



# The MINER $\nu$ A Experiment



Elaine Schulte

Rutgers, The State University of New Jersey

Piscataway, NJ

On Behalf of the  
MINER $\nu$ A Collaboration

# Introduction




1. What physics topics are accessible to Minerva?
2. Why are these topics important?
3. How will Minerva address this physics?

# Physics Topics Accessible to Minerva



<i>Physics Topic</i>	<i>Experimental Need</i>
<b>Deeply-Inelastic Scattering:</b>	<b>Improved statistics with better determination of final states</b>
<b>Quasi-Elastic Scattering: Axial Form Factor of Nucleon</b>	<b>Improved Precision over a wide <math>Q^2</math> Range</b>
<b>Coherent Scattering: Single Pion Production</b>	<b>1) Improved statistical precision of total cross section</b> <b>2) Measurements of nuclear dependence (A-dependence)</b>
<b>Resonance Production: Both Neutral Current (NC) and Charged Current (CC)</b>	<b>1) Improved statistical precision with 1-5 GeV neutrinos</b> <b>2) Quark-Hadron Duality</b>
<b>Nuclear Physics</b>	<b>Precision studies of neutrino-nucleus scattering as compared to charged lepton-nucleus scattering.</b>

# Existing Measurements of Neutrino-Nucleon Cross Section






# Quasi-Elastic Neutrino Scattering

$$\nu_{\mu}(\bar{\nu}_{\mu})n(p) \rightarrow \mu^{+}(\mu^{-})p(n)$$

$$\langle p | J_{\lambda}^{+} | n \rangle = \bar{u} \left[ \gamma_{\lambda} F_V^1(q^2) + i \sigma_{\lambda\nu} q^{\nu} \frac{\xi F_V^2(q^2)}{2M} + \gamma_{\lambda} \gamma_5 F_A(q^2) + q_{\lambda} \gamma_5 \frac{F_P(q^2)}{M} \right] u$$



$F_V^1(q^2)$  &  $F_V^2(q^2)$  are the Vector Form Factors

(extractable from  $G_E^N, G_M^N$ )

$F_A(q^2)$  is the Axial Form Factor


(extractable from neutrino scattering!)

# Form Factor Measurements



## MINER $\nu$ A Measurement of Axial FF

QE scattering,  $\nu_\mu$ ,  $F_A(Q^2)/\text{dipole}$ ,  $M_A=1.014$  GeV



Minerva estimated  $F_A(Q^2)$  statistical precision

based on Monte Carlo simulation attached to the electric form factor,  $G_E^p(Q^2)$ , for the nucleon to indicate scale.

The  $G_E^p(Q^2)$  scales used in this plot are based on polarization transfer measurements performed at Jefferson Lab (red) and measurements of the total elastic electron-nucleon scattering cross section (magenta).


**The Axial Form Factor of the Nucleon is poorly known...**




# Coherent Neutrino Scattering

$$\nu_{\mu} (\bar{\nu}_{\mu}) N \rightarrow \mu^+ (\mu^-) \pi N \text{ Charged Current}$$

$$\nu_{\mu} (\bar{\nu}_{\mu}) N \rightarrow \nu_{\mu} (\bar{\nu}_{\mu}) \pi N \text{ Neutral Current}$$




# Coherent Pion Production: A Window on the Weak Interaction



# Example of MINER $\nu$ A's Analysis Potential Coherent Pion Production



CC Coherent Pion Production Cross Section






# Resonance Production

$$\nu_{\mu}(\bar{\nu}_{\mu}) n(p) \rightarrow \mu^{+}(\mu^{-}) R \text{ Charged Current}$$

$$\nu_{\mu}(\bar{\nu}_{\mu}) n(p) \rightarrow \nu_{\mu}(\bar{\nu}_{\mu}) R \text{ Neutral Current}$$

Form Factors are needed to describe the N-Resonance transitions.

- electron scattering probes vector component of these form factors
- neutrino scattering will probe axial component



Cross sections, thus the form factors, for neutrino excitation of resonances are virtually unknown.

# Quark-Hadron Duality




**Quark-Hadron Duality:** The relationship between the DIS structure function  $F_2$  and the average resonance cross section as measured in electron scattering.

The cause of quark-hadron duality is not well known...

Neutrino scattering will help untangle this phenomena since neutrino interactions explicitly provide insight into flavor dependent behavior.

## Quark-hadron duality



Niculescu et al.,  
*Phys. Rev. Lett.* 85 (2000) 1182

$$\xi = 2x / (1 + \sqrt{1 + 4M^2x^2/Q^2})$$



# Nuclear Effects

The Past: Neutrino interactions were measured on heavy nuclei with low statistical precision; nuclear effects could be ignored...but...


The present: interactions are being measured with increasing precision; nuclear effects are now important...so...

The future: precision  $A$ -dependence studies must be performed!



# Nuclear Effects



## Predicted difference $\nu$ -A vs $e/\mu$ -A



MINER $\nu$ A should be able to determine this ratio to a few percent for  $n > 6$  GeV.



# Why are these Topics Important?



<i>Physics Topic</i>	<i>Experimental Need</i>
<b>Deeply-Inelastic Scattering:</b>	Observed energy in detectors can be obscured by final state interactions in nuclear media.
<b>Quasi-Elastic Scattering: Axial Form Factor of Nucleon</b>	<ol style="list-style-type: none"> <li>1) The axial form factor of the nucleon is poorly known.</li> <li>2) Cross section uncertainties are a major portion of oscillation experiment error budgets.</li> </ol>
<b>Coherent Scattering: Single Pion Production</b>	<ol style="list-style-type: none"> <li>1) Coherent scattering is, in general, a good probe of the weak interaction.</li> <li>2) Coherent scattering is expected to be a large background for future precision neutrino oscillation experiments.</li> </ol>
<b>Resonance Production: Both Neutral Current (NC) and Charged Current (CC)</b>	Improved understanding of the transition from quasi-elastic processes to deeply-inelastic scattering processes in the weak sector.
<b>Nuclear Physics</b>	<ol style="list-style-type: none"> <li>1) Nuclear medium dependence of neutrino interactions is important for interpretation of future neutrino oscillation studies.</li> <li>2) Differences are expected between charged and neutral lepton structure functions.</li> </ol>

# For Example: Helping MINOS and NO $\nu$ A/T2K



## Measurement of $\Delta m^2$ with MINOS:

**Needed:** detailed understanding of the relationship between the incoming neutrino energy and the visible energy in the detector



**From:** precision cross section measurements and neutrino-initiated nuclear reactions

## Measurement of $\sin^2 \theta_{13}$ with NO $\nu$ A:

**Needed:** absolute cross sections of signal & background reactions

**From:** precision cross section measurements

*see: D. A. Harris, et al., hep-ex/041005 for further info...*



# How will Minerva Achieve these Physics Goals?



- ✗ Lots of Neutrinos
  - ✓ Provided by the NuMI Beam at FNAL
    - ✓ approximately  $10^3$  times more intense than previously available beams!
  
- ✗ Massive Detector with:
  - ✓ Good Tracking Resolution
  - ✓ Good Momentum Resolution
  - ✓ Low Momentum Particle Detection Threshold
  - ✓ Particle Identification Capabilities
  
- ✗ Array of Nuclear Targets
  - ✓ Carbon
  - ✓ Iron
  - ✓ Lead

# What is Minerva?

## Main INjector ExpeRiment $\nu$ -A\*



MINER $\nu$ A is a compact, fully active neutrino detector designed to study neutrino-nucleus interactions with unprecedented detail.

\*Minerva, pictured above, was the Roman goddess of wisdom and technical skill.

# The MINERvA Collaboration



D. Drakoulakos, P. Stamoulis, G. Tzanakos, M. Zois  
*University of Athens, Greece*

D. Casper#, J. Dunmore, C. Regis, B. Ziemer  
*University of California, Irvine*

E. Paschos  
*University of Dortmund*

M. Andrews, D. Boehnlein, N. Grossman,  
D. A. Harris#, N. Grossman, J. Kilmer, M.  
Kostin, J.G. Morfin\*, A. Pla-Dalmau,  
P. Rubinov, P. Shanahan, P. Spentzouris  
*Fermi National Accelerator Laboratory*

I. Albayrak, M.E. Christy, C.E. Keppel, V. Tvaskis  
*Hampton University*

R. Burnstein, O. Kamaev, N. Solomey  
*Illinois Institute of Technology*

S. Kulagin  
*Institute for Nuclear Research, Russia*

I. Niculescu, G. Niculescu  
*James Madison University*

G. Blazey, M.A.C. Cummings, V. Rykalin  
*Northern Illinois University*

Collaboration of:

Particle, Nuclear, and Theoretical  
physicists

W.K. Brooks, A. Bruell, R. Ent, D. Gaskell,  
W. Melnitchouk, S. Wood  
*Jefferson Lab*

L. Aliaga, J.L. Bazo, A. Gago  
*Pontificia Universidad Catolica del Peru*

S. Boyd, S. Dytman, M.-S. Kim, D. Naples, V. Paolone  
*University of Pittsburgh*

A. Bodek, R. Bradford, H. Budd, J. Chvojka,  
P. de Barbaro, R. Flight, S. Manly, K. McFarland\*,  
J. Park, W. Sakumoto, J. Steinman  
*University of Rochester*

R. Gilman, C. Glasshausser, X. Jiang,  
G. Kumbartzki, R. Ransome#, E. Schulte  
*Rutgers University*

A. Chakravorty  
*Saint Xavier University*

D. Cherdack, H. Gallagher, T. Kafka, W.A. Mann,  
W. Oliver  
*Tufts University*


R. Ochoa, O. Pereyar, J. Solana  
*Universidad Nacional de Ingenieria*

J.K. Nelson#, R. Schneider, F.X. Yumiceva  
*The College of William and Mary*


\* Co-Spokespersons

# Members of the MINERvA Executive Committee


# Lots of Neutrinos-NuMI Beam Line



MINOS  
 MINER $\nu$ A



# The MINER $\nu$ A Detector




- Nuclear Targets:
  - 1/2 Ton each carbon, iron, and lead
- Active Target:
  - 5.78 Tons segmented scintillator planes
- Electromagnetic Calorimeter:
  - Interleaved lead sheet (0.2 cm thick) with segmented scintillator planes
- Hadronic Calorimeter:
  - Interleaved Iron sheet (2.54 cm thick) with segmented scintillator planes




# Active Detector Elements


Basic element: 1.7x3.3cm triangular strips.  
 1.2mm WLS fiber readout in center hole



## MINER $\nu$ A optical system



# Front View of Detector



# Event Rates:

## 13 Million Total CC events

### 4 - year run



**Fiducial Volume:**  
**3 tons Polystyrene,  $\approx 0.6$  t C,  $\approx 1/2$  t Fe and  $\approx 1/2$  t Pb**  
**Expected CC event samples:**  
**8.6 M  $\nu$  events in Polystyrene**  
**1.4 M  $\nu$  events in C**  
**1.4 M  $\nu$  events in Fe**  
**1.4 M  $\nu$  events in Pb**

<i>Charged-Current Physics Topic</i>	<i>Expected Statistics</i>
	<b>3 Tons of Polystyrene</b>
<b>Quasi-Elastic</b>	<b>0.8 M</b>
<b>Resonance</b>	<b>1.6 M</b>
<b>Transition: Resonance to DIS</b>	<b>2 M</b>
<b>DIS and Structure Functions</b>	<b>4.1 M</b>
<b>Coherent Pion Production</b>	<b>85 K CC/37 K NC</b>

# Current Status of MINER $\nu$ A



- MINER $\nu$ A is an established project and approved by FermiLab.
- Component research and development, and prototyping, are well underway at our member institutions.
- Current scheduling model indicates construction starting in Oct. 2006 and installation-finishing/commissioning-starting in early Fall 2008.

# Summary



## 1. What physics topics are accessible to Minerva?

- A. Minerva will provide improved precision neutrino-nucleus cross section measurements at neutrino energies from 1 to 15 GeV.
- B. Minerva will be able to investigate DIS, quasi-elastic, coherent, and resonance processes with precision much improved over most present neutrino cross section measurements.

## 2. Why are these topics important?

- A. Coherent processes comprise a significant source of background for neutrino future neutrino oscillation studies.
- B. Cross sections for resonance production in neutrino scattering are relatively unknown.
- C. Nuclear medium effects from neutrino interactions are expected to differ from their charged-lepton counterparts.
- D. Neutrino-nuclear effects have not been studied in high-mass targets.
- E. The axial form factor of the nucleon is poorly known.

## 3. How will Minerva address this physics?

- A. Minerva is a multi-ton detector designed specifically for precision cross section measurements.
- B. Minerva will make use of the high-intensity neutrino beam from NuMI.
- C. Minerva is outfitted with an array of nuclear targets for the express purpose of high-precision studies of nuclear medium effects in neutrino interactions.