Electron neutrino quasielastic scattering

and

Observation of neutral-current diffractive-like process

at MINERvA

Jeremy Wolcott
University of Rochester

Fermilab Joint Experimental-Theoretical Seminar
September 18, 2015

Overview

- Introduction and motivation
  - Significance of ν_e cross sections for ν oscillation
  - The MINERvA experiment
- ν_e CCQE (0π) in MINERvA
  - Selection and reconstruction of events
  - Cross section extraction
- NC diffractive-like background
  - Characterization of event sample
  - Discussion of significance
- Summary and conclusions

Introduction and motivation

The promise of ν_e appearance

ν_e appearance is a centerpiece in modern ν physics!

The promise of ν_e appearance

ν_e appearance oscillation experiments can help answer **fundamental questions** about leptons:



e.g.: How are the neutrino masses distributed?

⬇

(Are neutrinos their own antiparticles?)

The promise of ν_e appearance

ν_e appearance oscillation experiments can help answer **fundamental questions** about leptons:



e.g.: How are the neutrino masses distributed?

⬇

(Are neutrinos their own antiparticles?)



e.g.: Do leptons violate CP symmetry?

⬇

(Why is there more matter than antimatter in the universe?)

How cross sections figure into the story

Have:

(intrinsic ν_e contamination has separate spectrum)

$\Phi_{\nu_\mu}\,\sigma_{\nu_\mu}$

$\Phi_{\nu_e}\,\sigma_{\nu_e}$

Need model!

Need:

$\Phi_{\nu_\mu}\,\sigma_{\nu_e}$

(oscillated ν_es have ν_μ spectrum)



How cross sections figure into the story



Traditionally:

- Get Φ_{ν_μ} from near detector constraint
- Lean on *lepton universality* to work out

σ_{ν_e} from σ_{ν_μ}

Why worry?

Lepton universality has been extensively checked.

So what's the problem?



FNAL (1990)

Phys. Rev. D **41**: 2653

(and many others)

Why worry?

Lepton universality has been extensively checked.

So what's the problem?

It's that #@%&! nucleus.

- Detectors built from complex materials
- Substituting m_e for m_μ exposes different range of available kinematics ($m_\mu/m_e \sim 200$)
  - Parameterizations used in ν-N cross sections have uncertainties – different amounts of uncertainty integrated for different kinematic space (Day and McFarland, *PRD* **86** 053003)
  - Nuclear effects (correlations between nucleons) different in different kinematic regimes (e.g., Pandey *et al.*, *PRC* **92** 024606; Nieves *et al.*, *PRC* 83 045501)



A direct measurement quantifies how safe $m_\mu \rightarrow m_e$ is.

Existing ν_e cross section measurements



$\sigma_e(E_\nu)$:
Gargamelle (1978) on CF_3Br;
T2K (2014) on CH

Nucl. Phys. B133, 2015
Phys. Rev. Lett. 113, 241803



$d\sigma_e/dE_e$, $d\sigma_e/d\theta_e$,
$d\sigma_e/dQ^2$:
T2K (2014) on CH

Phys. Rev. Lett. 113, 241803

Existing ν_e cross section measurements



$\sigma_e(E_\nu)$:
Gargamelle (1978) on CF_3Br;
T2K (2014) on CH
Nucl. Phys. B133, 2015
Phys. Rev. Lett. 113, 241803



$d\sigma_e/dE_e$, $d\sigma_e/d\theta_e$,
$d\sigma_e/dQ^2$:
T2K (2014) on CH
Phys. Rev. Lett. 113, 241803

Difficult measurement...
Low stats ↔ large errors, no exclusive reactions.

Gargamelle: 244 events at ~90% purity
T2K: 315 events at ~65% purity

Experimental prospects



$$\pi^+ \rightarrow \nu_\mu \mu^+$$

NuMI is the highest-intensity ν_μ beam in the world...

Experimental prospects



$$\pi^+ \rightarrow \nu_\mu \mu^+$$
$$\hookrightarrow e^+ \nu_e \bar{\nu}_\mu$$

NuMI is the highest-intensity ν_μ beam in the world...

... and therefore the highest-intensity (accelerator energy) ν_e beam in the world too.

An on-axis cross section experiment could potentially have the statistics to perform the first exclusive cross section measurement!

Experimental prospects



Triangular inner detector scintillator strips





MINERvA is designed for cross sections
(central fiducial region is a
fine-grained scintillator tracker)
and is on-axis in NuMI.

Perfect!

ν_e CCQE at MINERvA

Signal definition



More energy transferred to nucleus



Choose to pursue the quasi-elastic (CCQE) channel:

- Evaluation of exclusive-state model:
  - <u>Important signal reaction for off-axis ν oscillation</u> that is well-represented at MINERvA energies
  - Direct comparison to MINERvA ν_μ measurement
- Easier final-state electron identification (less other confusion in event)

Signal definition

Signal is really quasielastic-like:

- One electron or positron (MINERvA isn't magnetized)
- Any number of nucleons (nuclear effects; FSI tricky – reduce model dependence)
- No other hadrons

(This strategy should seem familiar from other data sets – e.g., MiniBooNE)



Event display of simulated
~4 GeV ν_e interaction in MINERvA

Event selection

- Stages of event selection:
  1) Muon elimination
  2) Selection of electromagnetic shower-like particles
  3) Rejection of photons
  4) CCQE-like event selection





Event selection

- Stages of event selection:
  1) **Muon elimination**
  2) Selection of electromagnetic shower-like particles
  3) Rejection of photons
  4) CCQE-like event selection







Event selection

- Stages of event selection:
  1) Muon elimination
  2) Selection of electromagnetic shower-like particles
  3) Rejection of photons
  4) CCQE-like event selection







Isolating ν_e-like events: EM-like final state selection

1. What fraction of energy is deposited at the track end?



3. What is the track's mean dE/dx?

Train a multivariate classifier using these three characteristics of the energy deposition profile of the shower-like object (then reject MIP-like tracks that slip through)

Isolating ν_e-like events: EM-like final state selection



3. What is the track's mean dE/dx?

Train a multivariate classifier using these three characteristics of the energy deposition profile of the shower-like object (then reject MIP-like tracks that slip through)





Event selection

- Stages of event selection:
  1) Muon elimination
  2) Selection of electromagnetic shower-like particles
  3) **Rejection of photons**
  4) CCQE-like event selection





π^0 background



Isolating ν_e events: Photon rejection



The energy deposition pattern early in the track helps discriminate between photons (background) and electrons

Isolating ν_e events: Photon rejection



The energy deposition pattern early in the track helps discriminate between photons (background) and electrons

Isolating ν_e events: Photon rejection





The energy deposition pattern early in the track helps discriminate between photons (background) and electrons

Isolating ν_e events: Photon rejection





The energy deposition pattern early in the track helps discriminate between photons (background) and electrons

Isolating ν_e events: Photon rejection





"Wait. That's a pretty serious disagreement!"

Yep, it is. (Now imagine if we used a detector incapable of fine-grained dE/dx, like a Cherenkov one...)

I'll return to this point shortly.

The energy deposition pattern early in the track helps discriminate between photons (background) and electrons

Event selection

- Stages of event selection:
  1) Muon elimination
  2) Selection of electromagnetic shower-like particles
  3) Rejection of photons
  4) CCQE-like event selection





ν_e inelastic background



Isolating ν_e-like events:
Quasielastic-like topology selection

Anything not within a 7.5° electron cone or a vertex activity region of 30 cm radius or tracked as a proton is "extra energy."



Simulated ν_e CCQE



CCQE

Simulated ν_e deep inelastic scattering



Isolating ν_e-like events: Quasielastic-like topology selection

Anything not within a 7.5° electron cone or a vertex activity region of 30 cm radius or tracked as a proton is "extra energy."

Cut on $\Psi = \frac{E_{extra}}{E_{cone}}$



Simulated ν_e CCQE

Simulated ν_e deep inelastic scattering





(Actual cut is a function of E_{vis}. This plot illustrates cut near most probable value of E_{vis} = 1.25 GeV.)

Selected sample



MC sample is 52.2% ν_e CCQE

(80.1% ν_e from any channel)







Infer ν kinematics from lepton's

(use QE hypothesis + stationary target assumption)



Steps to a cross section



Steps to a cross section



Constraining backgrounds

Sideband 1: contains Michel $e^{\pm}$ (rich in inelastic ν_e)



Sideband 2: larger "extra energy" (rich in inelastics)





Select two sidebands rich in the major backgrounds...

Constraining backgrounds



Michel-match sideband



Extra energy sideband



… and examine the normalizations of two distributions in each sideband (one of them, candidate electron energy, shown here; candidate electron angle is also used).

Constraining backgrounds



Michel-match sideband



Extra energy sideband



… and examine the normalizations of two distributions in each sideband (one of them, candidate electron energy, shown here; candidate electron angle is also used).

Then, fit the normalizations of the "other ν_e" and "other NC π^0"+"CC ν_μ π^0" categories, using the four distributions simultaneously.

Constraining backgrounds





This is one of four (sideband, variable) combinations that are fitted simultaneously.
(Here: electron energy in "extra energy" sideband.)

Scale factors:
0.90 for "other ν_e", 1.11 for "other NC π^0"+"CC ν_μ π^0"

Unmodeled data excess







The (minimal) contribution in the signal region from the “mystery” process is subtracted along with the other backgrounds.
Will return to the question of its identity shortly.

Steps to a cross section



Unfold the *observable* quantities, using a Bayesian technique, to correct for the simulated resolutions

$$\left(\frac{d\sigma}{d\xi}\right)_i = \frac{1}{\Phi} \times \frac{1}{T_n} \times \frac{1}{(\Delta\xi)_i} \times \frac{\sum_j U_{ij}\left(N_j^{obs} - N_j^{bknd}\right)}{\epsilon_i}$$

(unsmearing matrix)

(selected sample)

(predicted background)

(flux)

(number of targets)

(bin width)

(efficiency)

Steps to a cross section

$$\left(\frac{d\sigma}{d\xi}\right)_i = \frac{1}{\Phi} \times \frac{1}{T_n} \times \frac{1}{(\Delta\xi)_i} \times \frac{\sum_j U_{ij}\left(N_j^{obs} - N_j^{bknd}\right)}{\epsilon_i}$$

(unsmearing matrix)

(selected sample)

(predicted background)

(efficiency)

(flux)

(number of targets)

(bin width)

Corrections for the predicted efficiency applied in each variable separately



Steps to a cross section

$$\left(\frac{d\sigma}{d\xi}\right)_i = \frac{1}{\Phi} \times \frac{1}{T_n} \times \frac{1}{(\Delta\xi)_i} \times \frac{\sum_j U_{ij}\left(N_j^{obs} - N_j^{bknd}\right)}{\epsilon_i}$$

(unsmearing matrix)

(selected sample)

(predicted background)

(efficiency)

(flux)

(number of targets)

(bin width)



The *a priori* flux is constrained using a separate *in situ* measurement of the neutrino-electron elastic scattering rate (also constrains this background)



Uncertainty summary



Includes energy scale estimated using the π^0 mass peak in a separate measurement; resolutions; other detector effects

Constrained as noted previously

Mostly enters in background subtraction (from GENIE 2.6.2)

cross sections





The result and the prediction from GENIE 2.6.2 are statistically consistent.





ν_e-ν_μ comparisons

Q^2 is the fundamental independent variable in the CCQE models.
We can compare $d\sigma/dQ^2$ to a previous measurement from MINERvA on ν_μ
to directly test the principle of lepton universality our models rely on.





Both the ν_μ and the ν_e cross sections
more or less agree with the model.

ν_e-ν_μ comparisons

Q^2 is the fundamental independent variable in the CCQE models.
We can compare $d\sigma/dQ^2$ to a previous measurement from MINERvA on ν_μ
to directly test the principle of lepton universality our models rely on.





For scale, "agreeing with the model" is somewhat closer than another well-known puzzle in CCQE scattering

ν_e-ν_μ comparisons

Q^2 is the fundamental independent variable in the CCQE models.
We can compare $d\sigma/dQ^2$ to a previous measurement from MINERvA on ν_μ to directly test the principle of lepton universality our models rely on.



And thus their ratio does as well.
Conclusion: using $\sigma_{\nu\mu}$ for $\sigma_{\nu e}$ is justified.

ν_e-ν_μ comparisons

Q^2 is the fundamental independent variable in the CCQE models.
We can compare $d\sigma/dQ^2$ to a previous measurement from MINERvA on ν_μ
to directly test the principle of lepton universality our models rely on.





p.s.... beware: don't read too much into the shape.
The shape is not significant when the correlations in the uncertainties are taken into account.

Investigating
the mystery process



Excess


Absolutely normalized (3.49 × 10^20 P.O.T.)
Statistical uncertainties only
Events
Signal region
"Excess region"
(12.5σ local stat-only significance; 3.1σ w/ systematics)
Min. 100mm dE/dx in first 500mm (MeV/cm)

Excess

- We believe our excess is due to NC diffractive scattering from Hydrogen
  - MINERvA tracker is hydrocarbon (lots of H)
  - No default model in GENIE
  - Event characteristics very similar... (next slide)



PCAC-based NC coherent production from *nuclei*

(analogous)



PCAC-based NC diffractive production from *H*

Excess

- Characteristics of excess match diffractive process very well:
  1) Two-photon π^0 shower
  2) Coherent-like scattering:
     - Forward kinematics
     - Very little other energy
  3) Visible proton energy
- Predominantly higher-energy showers



NC diffractive production from *H*



Excess

- Characteristics of excess:
  1) Two-photon π^0 shower
  2) Coherent-like scattering:
     - Forward kinematics
     - Very little other energy
  3) Visible proton energy
- Predominantly higher-energy showers



Could it just be more electrons?





Michel electron sideband is heavily dominated by ν_e.

Very well modeled (χ^2/n.d.f. = 63.5/50).

Unlikely to be an electron shower modeling problem.

Could it be extra nuclear activity?



MINERvA muon neutrino CCQE found evidence that sometimes more particles are produced at the vertex than the simulation predicts.

Does the excess stem from overlap between extra particles and the electron shower?

Could it be extra nuclear activity?



MINERvA muon neutrino CCQE found evidence that sometimes more particles are produced at the vertex than the simulation predicts.

Does the excess stem from overlap between extra particles and the electron shower?

(Simulated signal event)



Use a sliding techinque that looks for the *minimum* 100mm dE/dx in the first 500mm of cone. Designed to "step over" overlaps from nuclear activity

Could it be extra nuclear activity?



MINERvA muon neutrino CCQE found evidence that sometimes more particles are produced at the vertex than the simulation predicts.

Does the excess stem from overlap between extra particles and the electron shower?



Toy studies with extra protons of 0-200 MeV added randomly to 25% of ν_e CCQE events do not create a measurable excess.

Not likely due to extra particles in ν_e CCQE events.

Identifying the shower energy

6.9 GeV simulated γ



4.8 GeV simulated π^0

($\pi^0 \rightarrow \gamma\gamma$ 99% b.f.)



16 GeV simulated η

($\eta \rightarrow \gamma\gamma$ 40%, $\eta \rightarrow 3\pi^0$ 30% b.f.)



Challenge:

Can't reliably differentiate between single- and multi-photon showers on event-by-event basis

Identifying the shower energy

6.9 GeV simulated γ



4.8 GeV simulated π^0

($\pi^0 \rightarrow \gamma\gamma$ 99% b.f.)



16 GeV simulated η

($\eta \rightarrow \gamma\gamma$ 40%, $\eta \rightarrow \pi^0\pi^0\pi^0$ 30% b.f.)



Challenge:

Can't reliably differentiate between single- and multi-photon showers on event-by-event basis

Attempt to separate *statistically*

Expect more extra energy (larger Ψ) for multi-photon shower





Disfavors η

Identifying the shower energy

6.9 GeV simulated γ



4.8 GeV simulated π^0

($\pi^0 \rightarrow \gamma\gamma$ 99% b.f.)



16 GeV simulated η

($\eta \rightarrow \gamma\gamma$ 40%, $\eta \rightarrow \pi^0\pi^0\pi^0$ 30% b.f.)



Challenge:

Can't reliably differentiate between single- and multi-photon showers on event-by-event basis

Attempt to separate *statistically*

Expect energy to be distributed asymmetrically around cone axis for multi-photon shower





Disfavors γ

Identifying the shower energy

6.9 GeV simulated γ



4.8 GeV simulated π^0

($\pi^0 \rightarrow \gamma\gamma$ 99% b.f.)



16 GeV simulated η

($\eta \rightarrow \gamma\gamma$ 40%, $\eta \rightarrow \pi^0\pi^0\pi^0$ 30% b.f.)



Challenge:

Can't reliably differentiate between single- and multi-photon showers on event-by-event basis

Attempt to separate *statistically*



Expect shower to be wider on average for multi-photon shower



Disfavors γ

Identifying the shower energy

So:

- Ruled out electrons
- Ruled out single photons
- Ruled out heavier mass state decaying to neutrals (η)

Conclusion:

Excess most consistent with showers from π^0s

Excess

- Characteristics of excess:
  1) Two-photon π^0 shower
  2) **Coherent-like scattering:**
     - **Forward kinematics**
     - **Very little other energy**
  3) Visible proton energy
- Predominantly higher-energy showers



Kinematics



$E\theta^2$ approximately energy-independent measure of "forwardness"

↓

Small $E\theta^2 \Rightarrow$ more forward

Conclusion:

Excess coherent-like in kinematics

What else is in these events?

Simulated ν_μ deep inelastic scattering w/ π^0





Extra energy ratio provides roughly shower-energy-indendent measure of other energy in event

$$\Psi = \frac{E_{extra}}{E_{cone}}$$

What else is in these events?

Simulated ν_μ deep inelastic scattering w/ π^0



Vertex energy (+ vertex-anchored tracks)



What else is in these events?

Simulated ν_μ deep inelastic scattering w/ π^0



Vertex energy (+ vertex-anchored tracks)

Extra energy ratio provides roughly shower-energy-indendent measure of other energy in event

$$\Psi = \frac{E_{extra}}{E_{cone}}$$





Excess again consistent with NC coherent, in which a π⁰ is the *only* final-state particle that interacts in the detector.

So: very little non-shower energy.

Excess

- Characteristics of excess:
  1) Two-photon π^0 shower
  2) Coherent-like scattering:
     - Forward kinematics
     - Very little other energy
  3) **Visible proton energy**
- Predominantly higher-energy showers



What else is in these events?



Data event in excess region



Measure **“upstream in-line” energy**: energy inside cone along same axis as shower cone but backwards.

What else is in these events?







Data event in excess region

Excess prefers more upstream inline energy than coherent (but not *much* more)...

... and the distance to it is consistent with an exponential falloff, which is the pattern expected if photons traveling from UIE location and pair-producing stochastically





Excess

- Characteristics of excess:
  1) Two-photon π^0 shower
  2) Coherent-like scattering:
     - Forward kinematics
     - Very little other energy
  3) Visible proton energy
- **Predominantly higher-energy showers**



Kinematic range

Observation:

Excess shower energy spectrum significantly stronger than GENIE NC π^0 models



NC diffractive π^0 off H

- GENIE *does* have one (beta-quality) implementation:
  - Based on model from D. Rein (*NPB* **278**:61, 1986)
  - Not enabled by default
    - Not vetted like default models
    - Would double-count: global single-pi tune predates this model
  - Our testing seems to imply implementation may have some issues
- We tried it anyway to see how it compares

NC diffractive π^0 off H

- GENIE *does* have one (beta-quality) implementation:
  - Based on model from D. Rein (*NPB* **278**:61, 1986)
  - Not enabled by default
    - Not vetted like default models
    - Would double-count: global single-pi tune predates this model
  - Our testing seems to imply implementation may have some issues
- We tried it anyway to see how it compares



Behavior in *dE/dx* promising...

NC diffractive π^0 off H

- GENIE *does* have one (beta-quality) implementation:
  - Based on model from D. Rein (*NPB* **278**:61, 1986)
  - Not enabled by default
    - Not vetted like default models
    - Would double-count: global single-pi tune predates this model
  - Our testing seems to imply implementation may have some issues
- We tried it anyway to see how it compares





... but kinematics are rather different than those of the excess...

NC diffractive π^0 off H

- GENIE *does* have one (beta-quality) implementation:
  - Based on model from D. Rein (*NPB* **278**:61, 1986)
  - Not enabled by default
    - Not vetted like default models
    - Would double-count: global single-pi tune predates this model
  - Our testing seems to imply implementation may have some issues
- We tried it anyway to see how it compares
- Qualitative features are similar
- We conclude that the excess is likely due to NC diffractive production, but this model cannot quantitatively predict it





... and its protons have too much KE for our UIE distribution

Cross section comparisons

Process	Total cross section integrated over MINERvA flux ($\times 10^{-39}$ cm^2) / CH
MINERvA data excess	0.19 ± 0.02 (stat) ± 0.08 (sys)
GENIE NC diffractive (Rein)	1.6
GENIE NC diffractive, $E_\pi > 3$ GeV	0.10
GENIE NC coherent (Rein-Sehgal)	1.8
GENIE NC coherent, $E_\pi > 3$ GeV	0.16
GENIE NC inclusive	~120
GENIE CC inclusive	~400

(GENIE 2.6.2 used for model calculations)

Excess process roughly corresponds in size to channels that are ~1% of total cross section in MINERvA flux.

(But, then imagine you're looking for rare ν_e oscillation events and can't separate e from γ...)

Summary and conclusions

- ν_e appearance oscillation measurements may yield fundamental insights in the lepton family
- ν_e cross sections are a vital ingredient in oscillation results, but challenge means few direct measurements exist
- MINERvA measurement of ν_e CCQE-like cross section:
  - Is first-ever exclusive-process ν_e cross section
  - Agrees with ν_μ CCQE measurement from MINERvA, supporting lepton universality hypothesis
- Observation of unpredicted NC diffractive-like process underscores need for sustained investment in generators

Thank you on behalf of MINERvA!

2015 collaboration meeting

Overflow

Numbers

- 3.49×10^{20} POT of data
- 2105 selected events (CCQE-like):

  estimated 1090 signal events (52% purity)

  + 604 other electron neutrino events (sample is 80% ν_e)

  + 54 other electron final state events (sample is 83% electron final state)
- Scale factors:
  - ν_e CC inelastic: 0.89 ± 0.08
  - NC & CC incoherent: 1.06 ± 0.12
- 371 selected events (excess)

The abundance of ν_μ cross sections





ν_μ ν_e

There is a wealth of ν_μ cross section data available, by comparison...

Isolating ν_e events: Data reduction

Event pre-selection:

- One (or more) reconstructed track(s) (>95% of $e^{\pm}$ in fiducial region begin with track)
- No obvious muons (never ν_e):
  - No tracks exiting back of detector
  - No Michel electron (e from μ decay) candidates (also rejects pions: $\pi^{\pm} \rightarrow \mu^{\pm} \rightarrow e^{\pm}$)



Simulated background rejected by muon cuts



Event display of simulated ~4 GeV ν_e interaction in MINERvA

PID variable: endpoint energy fraction

5.6 GeV e-



470 MeV p+



PID variable: endpoint energy fraction





1. Divide the energy deposits into bins of 10 g/cm^2 of areal density.

PID variable: endpoint energy fraction



2. Correct the energy deposits for the calorimetry.
3. Determine the median of the energy deposits (excluding the last one).

PID variable: endpoint energy fraction



4. Endpoint energy fraction = $\frac{E_{last}}{E_{median}}$

PID variable: shower “width”

Merge MIP-like pairs like this one (two brightest strips are neighbors) into one pseudo-strip with the sum of their charge



PID variable: shower “width”

(the blue ones were merged)



x

z

strip number



For each plane:

Take standard deviation of illuminated strip numbers (after merging), weighted by charge, in this plane

Then use the median of those standard deviations to characterize the event's “width”

Benchmarking the PID







Samples were selected by other analyses in MINERvA:

- $\pi^{\pm}$ from $\Delta^{\pm}$ arXiv:1406.6415
- p^{+} in CCQE in Phys.Rev. D91, 071301
- γ from π^{0} in Phys.Lett. B749, 130













Photon rejection cut





“Extra energy” cut

Anything not within a 7.5° electron cone or a vertex activity region of 30 cm radius or tracked as a proton is “extra energy.”



Simulated ν_e CCQE

Cut on $\Psi = \frac{E_{extra}}{E_{cone}}$



After marginalizing over all E_{vis}. Cut illustrated is around most probable value of E_{vis} = 1.25 GeV.



Fitting backgrounds: effect in signal region





Data
ν_e CCQE-like
Other CC ν_e
ν + e elastic
NC Coh
Other NC π^0
CC ν_μ π^0
Ad hoc π^0
Other

Signal region after fitting











Background constraint: θ_e





Before constraint

Michel electron sideband

Extra energy sideband





After constraint

Background constraint: E_e



Before constraint



Michel electron sideband

Extra energy sideband



After constraint



Background-subtracted distributions









Migration matrices









Unfolded distributions









Efficiency estimates





Mean selection efficiency is 35.3%.





Uncertainties on cross sections









Flux prediction: ancestry



Kaon-parent flux is almost exactly 10% of ν_e flux between 0 and 10 GeV



A priori uncertainties

Flux prediction: ancestry



Kaon-parent flux is almost exactly 10% of ν_e flux between 0 and 10 GeV



ν_μ flux for comparison

Effect of flux constraint





Reduction of 5-10% in prediction,
and 5-10 percentage points in predicted uncertainty as well

Detector model uncertainties



Based on studies of track reconstruction From MINOS

Based on bench and *in situ* measurements



1.8% EM energy scale uncertainty from fitting π^0 mass peak

Based on destructive assays of detector planes

Interaction model uncertainties



Computed by using single photon model to fit excess instead of π^0

Statistical uncertainties in sideband affect background fitting

GENIE generator uncertainties are dominated by uncertainties on inelastic pion interactions and pion absorption in final-state interaction model (both of which affect the content of the background prediction and the prediction of the signal within the sidebands)

Effect of correlations in systematics



Variation in EM energy scale makes slope of data distribution much closer to agreeing with MC.

Brings net x^2 down.

Acceptance comparison





Nuisance distributions (1)







Nuisance distributions (2)





ν_e CCQE-like | Other CC ν_e | ν + e elastic | NC Coh

Other NC π^0 | CC ν_μ π^0 | Other

GENIE CCQE vs CCQE-like





Particle identification



An invaluable tool: **shape comparisons**

Particle identification





An invaluable tool: **shape comparisons**

Particle identification





An invaluable tool: **shape comparisons**



Compare data shape to model shapes

Ruling out externally-entering sources

If front-entering particles responsible, would expect excess to cluster around front of fiducial region.

No such behavior observed.







Ruling out externally-entering sources

Project onto radial coordinate ρ





If side-entering particles responsible, would expect excess to cluster around edges of fiducial region and point inwards.

No such behavior observed.







Isn't this just NC coherent?







Doubling coherent to fit excess would create strong tension with MINERvA CC coherent measurement (same model produces CC and NC coherent in GENIE)

Isn't this just NC coherent?



(total π^0 energy)



Doubling coherent to fit excess would create strong tension with NOMAD NC coherent measurement

No: not NC coherent.

GENIE NC diffractive model

The GENIE model produces roughly the expected proton kinematics, but the charged-current version significantly over-predicts compared to what data is available (*Nucl.Phys.B* **264** 221), unlike in Rein's paper







Excess region scan

Takeaways:

- Events scanned as "double shower" are always π^0s
- π^0s frequently appear as "single showers" in the sample

Interpretation:

- Data excess region has significantly **more π^0s** than MC
- Data excess region has **more in-line activity** than MC



support our algorithmic conclusions

Category clarifications:

- "+ vertex" means activity near the shower vertex
- "+ remote" means activity away from the shower
- "+ in-line" means activity away from the shower, but upstream, in-line with the shower axis